

On MT -Convexity

Mevlüt TUNÇ and Hüseyin YILDIRIM

ABSTRACT. In this paper, one new classes of convex functions which is called MT -convex functions are given. We also establish some Hadamard-type inequalities.

1. Introduction

The following definition is well known in the literature: A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if inequality

$$(1.1) \quad f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

holds for all $x, y \in I$ and $t \in [0, 1]$. Geometrically, this means that if P, Q and R are three distinct points on the graph of f with Q between P and R , then Q is on or below chord PR .

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function and $a, b \in I$ with $a < b$. The following double inequality:

$$(1.2) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

is known in the literature as Hadamard's inequality (or H-H inequality) for convex function. Keep in mind that some of the classical inequalities for means can come from (1.2) for convenient particular selections of the function f . If f is concave, this double inequality hold in the inversed way.

In [3], Pachpatte established two Hadamard-type inequalities for product of convex functions.

THEOREM 1. Let $f, g : [a, b] \subseteq \mathbb{R} \rightarrow [0, \infty)$ be convex functions on $[a, b]$, $a < b$. Then

$$(1.3) \quad \frac{1}{b-a} \int_a^b f(x)g(x)dx \leq \frac{1}{3}M(a, b) + \frac{1}{6}N(a, b)$$

and

$$2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)g(x)dx + \frac{1}{6}M(a, b) + \frac{1}{3}N(a, b)$$

2000 *Mathematics Subject Classification.* 26D15.

Key words and phrases. convexity, AM-GM inequality.

where $M(a, b) = f(a)g(a) + f(b)g(b)$ and $N(a, b) = f(a)g(b) + f(b)g(a)$.

We recall the well-known AM-GM inequality for n positive real numbers which can be stated as follows.

If $p_1, \dots, p_n$ are positive numbers which sum to 1 and f is a real continuous function that is concave up, then

$$\sum_{i=1}^n p_i f(x_i) \geq f\left(\sum_{i=1}^n p_i x_i\right)$$

A special case is

$$\sqrt[n]{x_1 x_2 \dots x_n} \leq \frac{x_1 + x_2 + \dots + x_n}{n}$$

with equality iff $x_1 = x_2 = \dots = x_n$.

We recall the well-known AM-GM inequality for two positive real numbers which can be stated as follows.

If $x, y \in \mathbb{R}^+$, then

$$(1.4) \quad \sqrt{xy} \leq \frac{x+y}{2}.$$

with equality if and only if $x = y$.

This inequality has many simple proofs. For example, a proof based on the concavity of the logarithmic function is presented in various sources, and the original reference is Jensen's paper [1]. A proof based on induction, given by Cauchy in 1821, is presented in many sources, as for example in [2], pp. 1–2.

In the following section our main results are given. We establish new a class of convex functions and then we obtain new Hadamard type inequalities for the new class of convex function.

DEFINITION 1. [See [4]] Two functions $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ are said to be similarly ordered, shortly f s.o. g , if

$$(f(x) - f(y))(g(x) - g(y)) \geq 0$$

for every $x, y \in X$.

2. MT-Convexity and Related Results

REMARK 1. If we take $x = t$ and $y = 1 - t$ in (1.4), we have

$$(2.1) \quad 1 \leq \frac{1}{2\sqrt{t(1-t)}}$$

for all $t \in (0, 1)$.

DEFINITION 2. A function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to belong to the class of $MT(I)$ if it is nonnegative and for all $x, y \in I$ and $t \in (0, 1)$ satisfies the inequality;

$$(2.2) \quad f(tx + (1-t)y) \leq \frac{\sqrt{t}}{2\sqrt{1-t}} f(x) + \frac{\sqrt{1-t}}{2\sqrt{t}} f(y).$$

REMARK 2. In (2.2), if we take $t = 1/2$, inequality (2.2) reduce to Jensen convex.

THEOREM 2. Let $f \in MT(I)$, $a, b \in I$ with $a < b$ and $f \in L_1[a, b]$. Then

$$(2.3) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx$$

and

$$(2.4) \quad \frac{2}{b-a} \int_a^b \tau(x) f(x) dx \leq \frac{f(a) + f(b)}{2}$$

where $\tau(x) = \frac{\sqrt{(b-x)(x-a)}}{b-a}$, $x \in [a, b]$

PROOF. Since $f \in MT(I)$, we have, for all, $x, y \in I$ (with $t = \frac{1}{2}$ in (2.2)) that

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2}$$

i.e. with $x = ta + (1-t)b$, $y = (1-t)a + tb$,

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{2} (f(ta + (1-t)b) + f((1-t)a + tb)).$$

By integrating, we get

$$(2.5) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{2} \left[\int_0^1 f(ta + (1-t)b) dt + \int_0^1 f((1-t)a + tb) dt \right],$$

Since

$$\int_0^1 f(ta + (1-t)b) dt = \int_0^1 f((1-t)a + tb) dt = \frac{1}{b-a} \int_a^b f(u) du,$$

we get the inequality (2.3) from (2.5).

For the proof of (2.4), we first note that if $f \in MT(I)$, then for all $a, b \in I$ and $t \in [0, 1]$, it yields

$$2\sqrt{t(1-t)}f(ta + (1-t)b) \leq tf(a) + (1-t)f(b)$$

and

$$2\sqrt{t(1-t)}f((1-t)a + tb) \leq (1-t)f(a) + tf(b).$$

By adding these inequalities and integrating on t over $[0, 1]$, we obtain

$$(2.6) \quad \int_0^1 \sqrt{t(1-t)} [f(ta + (1-t)b) + f((1-t)a + tb)] dt \leq \frac{f(a) + f(b)}{2}$$

Therefore,

$$(2.7) \quad \begin{aligned} \int_0^1 \sqrt{t(1-t)} f(ta + (1-t)b) dt &= \int_0^1 \sqrt{t(1-t)} f((1-t)a + tb) dt \\ &= \frac{1}{b-a} \int_0^1 \frac{\sqrt{(b-x)(x-a)}}{(b-a)^2} f(x) dx. \end{aligned}$$

We get (2.4) by combining (2.6) with (2.7) and the proof is completed.

The constant 1 in (2.3) is the best possible because this inequality obviously reduces to an equality for the function $f(x) = 1$ for all $a \leq x \leq b$. Additionally,

this function is to in the class $MT(I)$, because

$$\begin{aligned} \frac{\sqrt{t}}{2\sqrt{1-t}}f(x) + \frac{\sqrt{1-t}}{2\sqrt{t}}f(y) &\geq \frac{\sqrt{t}}{2\sqrt{1-t}} + \frac{\sqrt{1-t}}{2\sqrt{t}} = g(t) \\ &\geq \min g(t) = g\left(\frac{1}{2}\right) \quad 0 < t < 1 \\ &= 1 \geq f(tx + (1-t)y) \end{aligned}$$

for all $x, y \in [a, b]$ and $t \in [0, 1]$. Thus, the proof is completed. $\square$

REMARK 3. In (2.4), if we take $x = \frac{a+b}{2}$, inequality (2.4) reduce to Jensen's inequality.

THEOREM 3. Let $f \in MT(I)$, $a, b \in I$ with $a < b$ and $f \in L_1[a, b]$. Then

$$\frac{\pi}{2}f\left(\frac{a+b}{2}\right) \leq f(a) + f(b).$$

PROOF. Since $f \in MT(I)$, we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq f\left(\frac{ta + (1-t)b}{2} + \frac{(1-t)a + tb}{2}\right) \\ &\leq \frac{1}{2}(f(ta + (1-t)b) + f((1-t)a + tb)) \\ &\leq \frac{1}{2}\left(\frac{\sqrt{t}}{2\sqrt{1-t}} + \frac{\sqrt{1-t}}{2\sqrt{t}}\right)(f(a) + f(b)) \end{aligned}$$

Moreover

$$4\sqrt{t(1-t)}f\left(\frac{a+b}{2}\right) \leq f(a) + f(b)$$

By integrating, we get

$$\begin{aligned} 4f\left(\frac{a+b}{2}\right) \int_0^1 \sqrt{t(1-t)} dt &\leq f(a) + f(b) \\ \frac{\pi}{2}f\left(\frac{a+b}{2}\right) &\leq f(a) + f(b). \end{aligned}$$

The proof is completed. $\square$

THEOREM 4. Let $f, g \in MT(I)$, $a, b \in I$ with $a < b$ and $fg \in L_1[a, b]$. Then we have the inequality

$$\begin{aligned} (2.8) \quad &\frac{1}{b-a} \int_a^b \mu(x) f(x) g(x) dx \\ &\leq \frac{1}{12} [f(a)g(a) + f(b)g(b)] + \frac{1}{24} [f(a)g(b) + f(b)g(a)] \end{aligned}$$

where $\mu(x) = \frac{(b-x)(x-a)}{(b-a)^2}$, $x \in [a, b]$.

PROOF. Since $f, g \in MT(I)$, we have

$$\begin{aligned} f(ta + (1-t)b) &\leq \frac{\sqrt{t}}{2\sqrt{1-t}}f(a) + \frac{\sqrt{1-t}}{2\sqrt{t}}f(b) \\ g(ta + (1-t)b) &\leq \frac{\sqrt{t}}{2\sqrt{1-t}}g(a) + \frac{\sqrt{1-t}}{2\sqrt{t}}g(b) \end{aligned}$$

Since f and g are nonnegative, we write that

$$\begin{aligned} &f(ta + (1-t)b)g(ta + (1-t)b) \\ &\leq \frac{t}{4(1-t)}f(a)g(a) + \frac{1}{4}(f(a)g(b) + f(b)g(a)) + \frac{1-t}{4t}f(b)g(b) \\ &= \frac{t^2}{4t(1-t)}f(a)g(a) + \frac{t(1-t)}{4t(1-t)}(f(a)g(b) + f(b)g(a)) + \frac{(1-t)^2}{4t(1-t)}f(b)g(b) \end{aligned}$$

Consequently,

$$\begin{aligned} &t(1-t)f(ta + (1-t)b)g(ta + (1-t)b) \\ &\leq \frac{1}{4}\left\{t^2f(a)g(a) + t(1-t)[f(a)g(b) + f(b)g(a)] + (1-t)^2f(b)g(b)\right\} \end{aligned}$$

By integrating, we get

$$\begin{aligned} &\frac{1}{b-a} \int_a^b \frac{(b-x)(x-a)}{(b-a)^2} f(x)g(x) dx \\ &\leq \frac{1}{12}[f(a)g(a) + f(b)g(b)] + \frac{1}{24}[f(a)g(b) + f(b)g(a)]. \end{aligned}$$

The proof is completed. $\square$

REMARK 4. If we choose $x = \frac{a+b}{2}$ in the inequality (2.8), we obtain the special state of the inequality (1.3).

THEOREM 5. Let f, g be similarly ordered, nonnegative and MT -convex functions on I , $a, b \in I$ with $a < b$ and $fg \in L_1[a, b]$. Then we have the inequality

$$\frac{1}{b-a} \int_a^b \mu(x) f(x)g(x) dx \leq \frac{f(a)g(a) + f(b)g(b)}{8}$$

where $\mu(x) = \frac{(b-x)(x-a)}{(b-a)^2}$, $x \in [a, b]$.

PROOF. Since $f, g \in MT(I)$, we have

$$\begin{aligned} f(ta + (1-t)b) &\leq \frac{\sqrt{t}}{2\sqrt{1-t}}f(a) + \frac{\sqrt{1-t}}{2\sqrt{t}}f(b) \\ g(ta + (1-t)b) &\leq \frac{\sqrt{t}}{2\sqrt{1-t}}g(a) + \frac{\sqrt{1-t}}{2\sqrt{t}}g(b) \end{aligned}$$

Since f and g are nonnegative and similarly ordered, we write that

$$\begin{aligned}
& f(ta + (1-t)b)g(ta + (1-t)b) \\
& \leq \frac{t}{4(1-t)}f(a)g(a) + \frac{1}{4}(f(a)g(b) + f(b)g(a)) + \frac{1-t}{4t}f(b)g(b) \\
& = \frac{t^2}{4t(1-t)}f(a)g(a) + \frac{t(1-t)}{4t(1-t)}(f(a)g(b) + f(b)g(a)) + \frac{(1-t)^2}{4t(1-t)}f(b)g(b) \\
& \leq \frac{t^2}{4t(1-t)}f(a)g(a) + \frac{t(1-t)}{4t(1-t)}(f(a)g(a) + f(b)g(b)) + \frac{(1-t)^2}{4t(1-t)}f(b)g(b)
\end{aligned}$$

Consequently,

$$\begin{aligned}
& 4t(1-t)f(ta + (1-t)b)g(ta + (1-t)b) \\
& \leq t^2f(a)g(a) + t(1-t)[f(a)g(a) + f(b)g(b)] + (1-t)^2f(b)g(b) \\
& = tf(a)g(a) + (1-t)f(b)g(b)
\end{aligned}$$

By integrating, we get

$$\frac{4}{b-a} \int_a^b \mu(x)f(x)g(x)dx \leq \frac{1}{2}[f(a)g(a) + f(b)g(b)].$$

The proof is completed. $\square$

References

- [1] J.L.W.V. Jensen, Sur les fonctions convexes et les inéqualités entre les valeurs moyennes, Acta Mathematica 30 (1906), 175–193.
- [2] B. Bollobás, Linear Analysis, an introductory course (Cambridge Univ. Press 1990).
- [3] B.G. Pachpatte, On some inequalities for convex functions, RGMIA Res. Rep. Coll., 6 (E), 2003.
- [4] H.J. Skala, On the characterization of certain similarly ordered super-additive functionals, Proceedings of the American Mathematical Society, 126 (5) (1998), 1349-1353.

DEPARTMENT OF MATHEMATICS, FACULTY OF ART AND SCIENCES, KILIS 7 ARALIK UNIVERSITY, KILIS, 79000, TURKEY

E-mail address: mevluttunc@kilis.edu.tr

DEPARTMENT OF MATHEMATICS, FACULTY OF ART AND SCIENCES, KAHRAMANMARAŞ SÜTÇÜ İMAM UNIVERSITY, KAHRAMANMARAŞ, 46000, TURKEY

E-mail address: hyildir@ksu.edu.tr